

# CRUCIFERS (Cabbage, Cauliflower, Broccoli, Chinese Cabbage)



GROWTH STAGES	Seedbed		Transplanting	Vegetative				Heading to Head Development			Maturity
	1-15 DAS	16-25 DAS	1-7 DAT	8-14 DAT	15-21 DAT	22-28 DAT	29-35 DAT	36-42 DAT	43-49 DAT	50-56 DAT	57-70 DAT
INSECTS	Cutworms / Armyworms										
					Diamond-back Moths						
	Webworms										
	Cabbage Worms										
	Leafminers										
RECOMMENDED INSECTICIDES Use Rate / 16L water tank load	<b>Sumicidin</b> 5-8 Tbsp		<b>Padan</b> 1.5- 2.0 Tbsp	<b>Sumicidin</b> 5-8 Tbsp	<b>Padan</b> 1.5- 2.0 Tbsp	<b>Sumicidin</b> 5-8 Tbsp	<b>Padan</b> 1.5- 2.0 Tbsp	<b>Sumicidin</b> 5-8 Tbsp	<b>Padan</b> 1.5- 2.0 Tbsp	<b>Sumicidin</b> 5-8 Tbsp	
DISEASES	Downy Mildew										
	Alternaria Leaf Spot										
RECOMMENDED FUNGICIDE Use Rate / 16L water tank load	<b>Manzate® DF</b> 3-6 Tbsp										



Specialization	For Control of Slugs	For pre-plant	Soil Conditioner and Growth Agent	Plant Nutrition	For Farmers Who Practice Insecticide Tank - mixing	Contact Fungicide in Wettable Powder Formulation
Recommended Specialist	BAYONET Spot Application	SLASH 16-36* Tbsp	LIMELITE 40 g	OMEX BIO 20 11- 12 Tbsp	SUMICIDIN 5-8 Tbsp	MANZATE 200 3-6 Tbsp



# BEANS (Baguio beans, Sweetpea)



GROWTH STAGES	Seedbed	Vegetative		Flowering to Fruiting		Harvesting	
	1-15 DAS	16 - 22 DAS	23 - 30 DAS	31 - 37 DAS	38 - 45 DAS	46 - 90 DAS	
INSECTS	Beanflies						
	Cutworms/Armyworms/Ants						
	Leafminers						
	Leafrollers						
	Aphids/Leafhoppers				Pod Borers		
RECOMMENDED INSECTICIDES Use Rate / 16L water tank load	<b>LORSBAN 3E PLUS</b> 2.5 - 3.5 Tbsp					<b>LEGEND</b> 1.5-2 Tbsp	<b>CHIX</b> 1.6-3.2 Tbsp
DISEASES	Downy Mildew						
	Anthracnose						
	Bean Rust						
RECOMMENDED FUNGICIDE Use Rate / 16L water tank load	<b>Manzate® DF</b> 3-6 Tbsp						



Specialization	For pre-plant	Soil Conditioner and Growth Agent	Plant Nutrition	For Farmers Who Practice Insecticide Tank - mixing	Contact Fungicide in Wettable Powder Formulation
Recommended Specialist	SLASH 16-36* Tbsp	LIMELITE 40 g	OMEX BIO 20 11- 12 Tbsp	SUMICIDIN 5-8 Tbsp	MANZATE 200 3-6 Tbsp



# TOMATOES



GROWTH STAGES	Seedbed	Vegetative		Flowering to Fruiting			Harvesting	
	1-30 DAT	1-15 DAS	16-30 DAT	31-37 DAT	38-45 DAT	46-52 DAT	53-60 DAT	61-90 DAT
INSECTS	Cutworms/Ants							
	Aphids/ Thrips/ Whiteflies						Fruitworms	
RECOMMENDED INSECTICIDES Use Rate / 16L water tank load	<b>Sumithion</b> 3.0 - 5.0 Tbsp							<b>CHIX</b> 1.6-3.2 Tbsp
DISEASES	Anthracnose							
	Early and Late Blight							
	<b>DANJIRI</b> 4-6 Tbsp	<b>Manzate® DF</b> 3-6 Tbsp	<b>DANJIRI</b> 4-6 Tbsp	<b>Manzate® DF</b> 3-6 Tbsp	<b>DANJIRI</b> 4-6 Tbsp	<b>Manzate® DF</b> 3-6 Tbsp	<b>DANJIRI</b> 4-6 Tbsp	<b>Manzate® DF</b> 3-6 Tbsp



Specialization	For pre-plant weed clearing	Plant Nutrition	Contact Fungicide in Wettable Powder Formulation
Recommended Specialist	SLASH 16-36* Tbsp	OMEX CALMAX 8-10 Tbsp	MANZATE 200 3-6 Tbsp

